

Relaxation of the Phase In an Inhomogeneous Bose Gas

V. S. Babichenko

R.S.C. "Kurchatov Institute" Kurchatov sq. 46, Moscow 123182, Russia.

An effective action is obtained of a Bose gas in the bulk separated into two regions by a strong external potential depending on the single coordinate. The main attention is focused on the relaxation of the difference between phases of the weakly coupling condensates of the different bulk domains separated from each other by the external potential.

The experimental realization of the Bose-Einstein condensation at ultralow temperatures in atomic vapors [1] provides an example of the systems in which the approximation of the weakly non-ideal gas is well applicable. This fact is connected with the small density of particles in the systems concerned. The realization of such systems gives the possibility of the experimental investigations of the macroscopic quantum phenomenons of different types. Recently, the manifestations of the macroscopic quantum phase and its behavior are studied actively [1-8], [11].

The investigation of the kinetic phenomena due to the relaxation of the order parameter is of a doubtless interest for the study of macroscopic quantum phenomena. In the present work we study the spatial relaxation of the phase of the order parameter in the inhomogeneous Bose gas with the weak coupling between different spatial regions due to a barrier tunneling.

The system concerned in the present work is a Bose gas of the small density in the bulk separated into two regions by a strong external potential $U(\vec{r})$ which depends on the x coordinate and does not depend on $\vec{r}_\perp = (y, z)$, i.e., $U(\vec{r}) = U(x)$. For simplicity, we assume that the external potential has a rectangular shape $U(x) = U_0$ for $-d < x < d$ and $U(x) = 0$ beyond this region of the x coordinate. The height of the external potential U_0 is supposed to be the largest energy parameter in the system, in particular, $U_0 \gg \mu$ where μ is the chemical potential $\mu = n\lambda$, λ is the scattering amplitude of Bose particles, and n is the density of the Bose gas. Due to this assumption the interaction between particles of the Bose gas in the region of the influence of the external potential, i.e., in the region $-d < x < d$, can be neglected.

The left and the right domains of the bulk are supposed to have the same temperature T smaller than the Bose-condensation temperature T_c . For simplicity, we suppose that in the initial state the density of the left and the right domains of the bulk are the same, but the phases of the left and the right Bose condensates are different. This assumption means that at the initial time moment there is nonzero current from one side of the bulk to another. This non-equilibrium initial state will relax to the equilibrium one which has the same densities and the same phases of the condensates for both sides of the bulk. This relaxation process is studied in the present work.

Note that in the case of superconductors the similar initial non-equilibrium state results in the Josephson oscillations with a small damping [9], [10]. The essential distinction of the Bose gas consisting of neutral atoms from superconductors in which the Cooper pairs represent the charged objects is the presence of the gapless excitation spectrum. The presence of low energy excitations results in the change of the character of the initial state relaxation making the relaxation essentially faster.

In the case of the homogeneous in y - z plane difference between the phases of the condensates the excitations radiated during the relaxation process have the one-dimensional character. The one-dimensional character of the radiated excitations results in a divergency of the small momentum correlators of these excitations. Hence, the consideration of the relaxation process to second order in the tunneling amplitude is not correct in contrast to the case of superconductors [10] and requires more accurate analysis.

I. PERTURBATION THEORY IN THE TUNNELING AMPLITUDE

As the introduction, we consider the perturbation theory in the amplitude of the tunneling under the external potential barrier. The Hamiltonian of the Bose gas with the tunneling term is

$$\hat{H} = \int_{U(x)=0} d^3r \left\{ \hat{\psi}^+(\vec{r}) \left(-\frac{1}{2m} \nabla^2 - \mu \right) \hat{\psi}(\vec{r}) + \frac{\lambda}{2} \left(\hat{\psi}^+(\vec{r}) \hat{\psi}(\vec{r}) \right)^2 \right\} + \quad (1.1)$$

$$+ \int d^3r_1 d^3r_2 \left\{ \hat{\psi}^+(\vec{r}_1) \hat{T}_{\vec{r}_1 \vec{r}_2} \hat{\psi}(\vec{r}_2) + \hat{\psi}^+(\vec{r}_2) \hat{T}_{\vec{r}_1 \vec{r}_2}^* \hat{\psi}(\vec{r}_1) \right\}$$

The form of the kernel of the integral operator $T_{\vec{r}_1 \vec{r}_2}$ in the case of the external potential independent of $\vec{r}_\perp$ can be taken as

$$T_{\vec{r}_1 \vec{r}_2} = \gamma_{x_1, x_2} \delta(\vec{r}_{\perp 1} - \vec{r}_{\perp 2}) \quad \text{where } x_1 > 0 \quad \text{and} \quad x_2 < 0 \quad (1.2)$$

The kernel γ_{x_1, x_2} is localized near $x_1 = x_2 = 0$ within the coordinate x of about a where $a \lesssim \xi$, ξ being the correlation length of the weakly interacting Bose gas $\xi = (n\lambda)^{-1/2}$. In the case of the slow variation of the operator $\hat{\psi}(\vec{r})$ compared with the scale ξ the kernel γ_{x_1, x_2} can be taken in the form

$$\gamma_{x_1, x_2} = \gamma_0 \delta(x_1) \delta(x_2) \quad (1.3)$$

The constant γ_0 is the small quantity due to the large magnitude of the external potential and obeys the inequality $\gamma_0 \ll \mu\xi$. The tunneling amplitude (1.3) will be obtained below and has the form $\gamma_0 = \frac{1}{2m} n \xi (\kappa \sinh(2\kappa d))^{-1}$ where $\kappa = \sqrt{2mU_0}$. Form (1.3) of the tunneling amplitude differs from that used in the work [11]. This difference is due to the behavior of the Fourier components of the tunneling amplitude $\gamma_{k_{1x}, k_{2x}}$ for the small momenta k_{1x}, k_{2x} . In contrast to the present work, in the work [11] the tunneling amplitude $\gamma_{k_{1x}, k_{2x}}$ vanishes $\gamma_{k_{1x}, k_{2x}} \rightarrow 0$ provided that the momenta k_{1x}, k_{2x} tends to zero $k_{1x}, k_{2x} \rightarrow 0$, this amplitude being nonzero for the condensate-condensate tunneling. This distinction has the far-reaching consequences, namely, nonzero value of $\gamma_{k_{1x}, k_{2x}}$ for $k_{1x}, k_{2x} \rightarrow 0$ results in the divergency of lower orders of the perturbation theory in the tunneling amplitude. This problem is discussed in detail below.

The characteristic frequency of the phase difference evolution between the right and left condensates is proportional to the small tunneling amplitude γ_0 . The characteristic momenta of particles corresponding to this frequency are small compared with the scale of the order of $1/\xi$. This means that the slow relaxation process of the phase difference is described by the slow variation in the spatial and temporal Fourier components of operators $\hat{\psi}_\alpha(\vec{r}, t)$. Index α is equal to $\alpha = 1, 2$ where 1,2 denotes the right-hand ($x > 0$) and the left-hand ($x < 0$) sides of the bulk, respectively. Therefore, the model form of the kernel γ_{x_1, x_2} given by Eq. (1.3) corresponds to the purpose of the work, namely, to describe the relaxation of the phase difference.

Due to the existence of the condensates in the left-hand and the right-hand sides of the bulk the operators $\hat{\psi}_\alpha(\vec{r}, t)$ can be represented in the well-known form

$$\hat{\psi}_\alpha(\vec{r}, t) = \Psi_\alpha(t) + \hat{\chi}_\alpha(\vec{r}, t) \quad (1.4)$$

$$\Psi_\alpha(t) = \sqrt{n_0} e^{i\varphi_\alpha(t)} \quad (1.5)$$

The fields $\Psi_\alpha(t)$ are the condensate fields for the right-hand ($\alpha = 1$) and the left-hand ($\alpha = 2$) sides of the bulk. The value n_0 is the Bose condensate density and $\varphi_\alpha(t)$ are the phases of the right-hand ($\alpha = 1$) and the left-hand ($\alpha = 2$) Bose-condensates. The operators $\hat{\chi}_\alpha(\vec{r}, t)$ are the annihilation operators of non-condensate particles. As has been written above, the main problem is the investigation of the time evolution of the magnitude $\Delta\varphi(t) = \varphi_1(t) - \varphi_2(t)$. For this purpose, we will obtain the effective action for the phases $\varphi_1(t)$ and $\varphi_2(t)$.

At first, we consider the perturbation theory in the tunneling amplitude for the small phase difference $\Delta\varphi(t)$. The diagrams for the renormalized tunnelling amplitude of the condensate particles have the form of the vertex with two external lines (as for the self-energy part) corresponding to the condensate field. The quantities corresponding to the incoming and outgoing dotted lines are Ψ_α and $\bar{\Psi}_\alpha$, respectively. To analyse the relaxation process, we use the Keldysh diagram technique with the Schwinger-Keldysh two-time contour [14], [15].

The reason of the impossibility to consider the lower orders of the perturbation theory in the tunneling amplitude is the divergency of the lower order diagrams when the external frequency ω tends to zero $\omega \rightarrow 0$. To demonstrate this, we consider the frequency Fourier transformation of one of second-order tunneling vertex

$$\Gamma_{11}^{(2)} = \gamma_0 g(\omega) \gamma_0 \quad (1.6)$$

where the function $g(\omega)$ denotes the Green function of the non-condensate fields [13] with the momentum $\vec{k}_\perp = 0$ and the same x -coordinates

$$g(\omega) = G_1\left(\omega, \vec{k}_\perp = 0; x_1 = x_2 = d\right) = G_2\left(\omega, \vec{k}_\perp = 0; x_1 = x_2 = -d\right) \quad (1.7)$$

The Green functions $G_\alpha(t - t', \vec{r} - \vec{r}')$ ($\alpha = 1, 2$) are the Belyaev Green functions of the non-condensate particles of the homogeneous interacting Bose gas. The momentum $\vec{k}_\perp$ corresponding to the Green function $g(\omega)$ vanishes $\vec{k}_\perp = 0$ due to the homogeneity of the condensate field in the y and z-directions. The Green functions $G_\alpha(\omega, \vec{k})$ and, correspondingly, $g(\omega)$ in the Keldysh diagrammatic technique has three components in triangle representation, namely, retarded $G^{(R)}$, advanced $G^{(A)}$ and kinetic $G^{(K)}$ Green functions [14], [15]. In the case of small frequencies $\omega \ll \mu$ the retarded Green function $g^{(R)}(\omega)$, for example, can be written in the form

$$g^{(R)}(\omega) = \int \frac{dp_x}{2\pi} \left(\frac{\mu}{(\omega + i\delta)^2 - c^2 p_x^2} \right) \quad (1.8)$$

The small momentum asymptotic of G in Eq. (1.8) is used due to the smallness of the external frequency ω ($\omega \ll \mu$). The main contribution into integral (1.8) is provided by the momenta $p_x \sim \omega/c \ll \mu/c$. Note that integral (1.8) has a singularity and, therefore, the magnitude of this integral is determined by the bypass around the pole. The calculation of the integral (1.8) gives

$$g^{(R)}(\omega) = -i \frac{\mu}{c} \frac{1}{2\omega} \quad (1.9)$$

This behavior of the function $g^{(R)}(\omega)$ means that for the sufficiently small magnitudes of ω , the consideration of some lower orders of the perturbation theory is not correct and the analysis of diagrams in all orders is necessary.

The diagrams of odd order of this perturbation theory results in the renormalization of the bare tunneling amplitude γ_0 . The sum of the odd order diagrams having one incoming and one outgoing line is denoted by the vertex $\Gamma_{\alpha\beta}$ ($\alpha \neq \beta$) where, as above, the indices $\alpha, \beta = 1, 2$ denote the right-hand or the left-hand sides of the bulk. The sum of the odd order diagrams having two incoming external lines or two outgoing external lines is denoted by the vertex $\hat{\Gamma}_{\alpha\beta}$ ($\alpha \neq \beta$) according to the notations of the self-energy part in the Belyaev diagram technique for the Bose systems [12], [13]. The term of the effective action corresponding to this renormalized tunneling amplitude has the form

$$S_{tunn} = \oint \oint dt dt' \bar{\Psi}_\alpha(t) \Gamma_{\alpha\beta}(t, t') \Psi_\beta(t') + \\ + \oint \oint dt dt' \left\{ \Psi_\alpha(t) \hat{\Gamma}_{\alpha\beta}(t, t') \Psi_\beta(t') + \bar{\Psi}_\alpha(t) \hat{\Gamma}_{\alpha\beta}(t, t') \bar{\Psi}_\beta(t') \right\}$$

The time contour in this expression is the Schwinger-Keldysh two-time contour. Note, that in the even orders of the tunneling amplitude the renormalized vertex $\Gamma_{\alpha\beta}$ has the coincident external indexes $\alpha = \beta$. These vertexes give the contribution to the self-energy part of the particles in one of the two subvolumes, in particular, the imaginary parts of these vertexes describe the relaxation process.

The renormalized vertexes connected with the dressed tunneling amplitude can be expressed by means of the Green functions taking into account the tunneling

$$\Gamma_{11} = \gamma_0 G_{22} \gamma_0 \quad (1.10)$$

$$\hat{\Gamma}_{11} = \gamma_0 F_{22} \gamma_0$$

$$\Gamma_{12} = \gamma_0 + \gamma_0 G_{21} \gamma_0$$

$$\hat{\Gamma}_{12} = \gamma_0 F_{21} \gamma_0$$

The system of the Dyson equations for $G_{\alpha\beta}$ and $F_{\alpha\beta}$ can be written as

$$G_{22} = G_2 + G_2 \gamma_0 G_{12} + F_2 \gamma_0 F_{12} \quad (1.11)$$

$$F_{22} = F_2 + F_2 \gamma_0 G_{12} + \tilde{G}_2 \gamma_0 F_{12}$$

$$G_{12} = G_1 \gamma_0 G_{22} + F_1 \gamma_0 F_{22}$$

$$F_{12} = \tilde{G}_1 \gamma_0 F_{22} + F_1 \gamma_0 G_{22}$$

Here G_α and F_α are the Belyaev Green functions of non-condensate particles of the homogeneous interacting Bose gas. The Green function G_α has one incoming and one outgoing end and the Green function F_α has two outgoing (or two incoming) ends. The Green function $\tilde{G}_\alpha$ denotes the Green function G_α with the opposite frequency ω and momentum $\vec{p}$, i.e., $\tilde{G}_\alpha(\omega, \vec{p}) = G_\alpha(-\omega, -\vec{p})$. This system of equations is written in the assumption of the small magnitude of the difference between phases of the left-hand and right-hand condensates. In the case of the small momenta, which is the most interesting for us, the Green functions G_α and F_α obeys the equality $G_\alpha = -F_\alpha$. In the case of the small momenta the solutions of Eqs. (1.11) can be simplified and the tunneling vertexes take the form

$$\Gamma_{11} = \frac{g(\omega) \gamma_0^2}{1 - 4\gamma_0^2 g^2(\omega)} \quad (1.12)$$

$$\hat{\Gamma}_{11} = -\frac{g(\omega) \gamma_0^2}{1 - 4\gamma_0^2 g^2(\omega)}$$

$$\Gamma_{12} = \gamma_0 + \frac{2g^2(\omega) \gamma_0^3}{1 - 4\gamma_0^2 g^2(\omega)}$$

$$\hat{\Gamma}_{12} = \frac{-2g^2(\omega) \gamma_0^3}{1 - 4\gamma_0^2 g^2(\omega)}$$

In the limit $\omega \rightarrow 0$ these vertexes can be written as

$$\Gamma_{11}^{(R)} \rightarrow -\frac{1}{4g(\omega)} = -i \frac{c}{2\mu} \omega \quad (1.13)$$

$$\hat{\Gamma}_{11}^{(R)} \rightarrow \frac{1}{4g(\omega)} = i \frac{c}{2\mu} \omega$$

$$\Gamma_{12}^{(R)} \rightarrow \frac{1}{2} \gamma_0$$

$$\hat{\Gamma}_{12}^{(R)} \rightarrow \frac{1}{2} \gamma_0$$

Note that the bare tunneling amplitude γ_0 is eliminated from the renormalized vertexes Γ_{11} , $\hat{\Gamma}_{11}$, and just these vertexes describe the relaxation of the phase (difference). This fact results in the large value of the relaxation rate.

The more rigorous and detailed consideration of the problem in the case of an arbitrary phase difference for the infinitely large size of the system in the x-direction is represented below. The assumption about the size of the system is accepted to focus the attention on the consideration of the relaxation process.

II. THE TUNNELING ACTION

Below the approach to the problem of the phase difference relaxation without using the perturbation theory in the tunneling amplitude is developed. For this purpose the effective action for the boundary values of the Bose field phases is obtained by the integration over the volume components of these fields with the fixed values of them at the left-hand and the right-hand sides of the external potential boundary. Integrating over these fields we take into account the inhomogeneous behavior of the condensate fields in the left-hand and the right-hand sides of the volume. The necessity of the consideration of the inhomogeneous condensate behavior even for the small momentums of the excitations is conditioned by the fact that the self consistent inhomogeneous condensate additional field makes the situation in some sense similar to the case of the scattering on the external potential having the discrete energy level which is very close to zero. This can results, for example, in the transmission coefficient of the order of unit including the case of the very small compared with μ energies [16]. The obtained effective action contains the relaxation part which is proportional to the first power of the frequency of the phase. For the small frequencies the relaxation part is

found to be much larger than the usual kinetic part of the phase dynamics which is proportional to the second power of the frequency. The consideration is developed in the framework of the Schwinger-Keldysh technique with the time reversing contour [14], [15].

The generating functional of the system can be written in the form

$$Z = \int D\psi D\bar{\psi} e^{i(S+S_j)} \quad (2.1)$$

The action S of the inhomogeneous non-equilibrium Bose gas is given by

$$S = \oint dt \int d^3r \left\{ \bar{\psi} [i\partial_t + \mu - U(x)] \psi - \frac{1}{2} (\bar{\nabla} \bar{\psi}) (\bar{\nabla} \psi) - \frac{\lambda}{2} (\bar{\psi} \psi)^2 \right\} \quad (2.2)$$

and term S_j reads as

$$S_j = \oint dt \int d^3r (\bar{\psi} j + \bar{j} \psi) \quad (2.3)$$

Here j and $\bar{j}$ are the infinitely small sources. The Planck constant $\hbar$ and mass m of a Bose particle are set equal to unity $\hbar = m = 1$.

Due to the large magnitude of potential U_0 in the region $x \in [-d, d]$ the interaction between Bose particles can be neglected and the integral for Z over the fields $\psi, \bar{\psi}$ takes the Gaussian form in this region of the x -coordinate. In this connection the integral for Z over the fields $\psi(x, \vec{r}_\perp; t), \bar{\psi}(x, \vec{r}_\perp; t)$ for $x \in [-d, d]$ with the fixed magnitudes of these fields at the boundary of the region of the nonzero external potential can be calculated. Moreover, the characteristic scale of the ψ -field variation in the time and in the $\vec{r}_\perp$ space is supposed to be much larger than $1/U_0$ and $1/\sqrt{U_0}$, respectively. It is these fields that describe the slow relaxation process possessing the characteristic frequency proportional to the small tunneling coefficient. Due to this assumption the tunneling amplitude of the ψ fields has the local character for the time variable t and space variables $\vec{r}_\perp$.

The magnitudes of the ψ -fields at the boundary of the region of the nonzero external potential are denoted as

$$\begin{aligned} \psi_1(x = d, \vec{r}_\perp; t) &= \psi_{s1}(\vec{r}_\perp; t) \\ \psi_2(x = -d, \vec{r}_\perp; t) &= \psi_{s2}(\vec{r}_\perp; t) \\ \bar{\psi}_1(x = d, \vec{r}_\perp; t) &= \bar{\psi}_{s1}(\vec{r}_\perp; t) \\ \bar{\psi}_2(x = -d, \vec{r}_\perp; t) &= \bar{\psi}_{s2}(\vec{r}_\perp; t) \end{aligned} \quad (2.4)$$

The calculation of the integral for Z over the ψ -fields in the region $x \in [-d, d]$ gives

$$Z = \int D\psi D\bar{\psi} e^{i(S_{vol} + S_{surf} + S_j)} \quad (2.5)$$

The part of the action S_{vol} is

$$S_{vol} = \oint dt \int_{U(x)=0} dx \int d^2r_\perp \left\{ \bar{\psi} (i\partial_t + \mu) \psi - \frac{1}{2} (\bar{\nabla} \bar{\psi}) (\bar{\nabla} \psi) - \frac{\lambda}{2} (\bar{\psi} \psi)^2 \right\} \quad (2.6)$$

where the region of the integration over the x -coordinate is the sum of the regions $-L < x < -d$ and $d < x < L$.

The part of the action S_{surf} can be represented as a sum of two terms $S_{surf}^{(\rho)}$ and $S_{surf}^{(tunn)}$ [17]

$$S_{surf} = S_{surf}^{(\rho)} + S_{surf}^{(tunn)} \quad (2.7)$$

$$S_{surf}^{(\rho)} = -\frac{\varkappa}{2} \coth(2\varkappa d) \oint dt \int d^2r_\perp \{ |\psi_{s1}(\vec{r}_\perp; t)|^2 + |\psi_{s2}(\vec{r}_\perp; t)|^2 \} \quad (2.8)$$

$$S_{surf}^{(tunn)} = \frac{\varkappa}{2 \sinh(2\varkappa d)} \oint dt \int d^2r_\perp \{ \bar{\psi}_{s1}(\vec{r}_\perp; t) \psi_{s2}(\vec{r}_\perp; t) + \bar{\psi}_{s2}(\vec{r}_\perp; t) \psi_{s1}(\vec{r}_\perp; t) \} \quad (2.9)$$

where the magnitude $\varkappa$ is $\varkappa = \sqrt{2U_0}$. Later on we assume that the width d of the potential barrier is sufficiently large so that the inequality $\varkappa d \gg 1$ takes place and, thus, $\sinh(2\varkappa d) \approx \cosh(2\varkappa d) \approx \frac{1}{2}e^{2\varkappa d} \gg 1$. The term $S_{surf}^{(tunn)}$ describes the tunneling between the right-hand and left-hand sides of the bulk. From the form of this tunneling part of the action it can easily be seen that the tunneling amplitude for the condensate particles and for the low energy non-condensate particles has the same nonzero magnitude in contrast with [11].

Our goal is the calculation of the functional integral for Z over the ψ -fields in the bulk, i.e., in the region $x \in (-L, -d) \cup (d, L)$ with the fixed values at the boundary of the region of the nonzero external potential, and, thus, obtaining the effective action for the fields at the boundary ψ_{s1}, ψ_{s2} .

III. THE SADDLE-POINT APPROXIMATION

Later on, it is convenient to change the variables in the following way

$$\begin{aligned} \vec{r} &\rightarrow \xi \vec{r} \\ t &\rightarrow \frac{1}{\mu} t \end{aligned} \quad (3.1)$$

where $\xi = 1/\sqrt{\mu}$. Thus, the system of units which we use later measures the quantities of the dimension of length in units of ξ and the quantities of the dimension of energy in units of μ .

Moreover, it is convenient to represent the fields ψ in the modulus-phase form

$$\psi = \rho e^{i\varphi}; \quad \bar{\psi} = \rho e^{-i\varphi} \quad (3.2)$$

The integral over the fields ψ in the right-hand and the left-hand sides of the bulk can be calculated within the framework of the saddle-point approximation. The saddle-point magnitude $\psi_{cl}(x, \vec{r}_\perp; t)$ of the field ψ obeys the Gross-Pitaevskii equation with the boundary conditions

$$\begin{aligned} \psi_{cl}(x = d, \vec{r}_\perp; t) &= \psi_{s1}(\vec{r}_\perp; t) \\ \psi_{cl}(x = -d, \vec{r}_\perp; t) &= \psi_{s2}(\vec{r}_\perp; t) \end{aligned} \quad (3.3)$$

In the present consideration we use the saddle-point approximation with respect to the modulus of the field ψ , i.e., ρ . The fluctuations of the modulus field ρ in the vicinity of the saddle-point magnitude $\rho^{(cl)}$ are taken into account within the framework of the perturbation theory. At the same time, the phase φ is considered exactly. The reason of such consideration is connected with the one-dimensional character of the excitations which are radiated during the relaxation of the difference between the phases of the left-hand and right-hand condensates. The saddle-point magnitudes $\rho^{(cl)}$ of the fields ρ have the form

$$\begin{aligned} \rho_1^{(cl)}(x, \vec{r}_\perp; t) &= \sqrt{n_0} F_1(x, \vec{r}_\perp; t) & (x > d) \\ \rho_2^{(cl)}(x, \vec{r}_\perp; t) &= \sqrt{n_0} F_2(x, \vec{r}_\perp; t) & (x < -d) \end{aligned} \quad (3.4)$$

where n_0 is the condensate density in the bulk far from the region of the nonzero external potential and

$$\begin{aligned} F_1(x, \vec{r}_\perp; t) &= \tanh(x - d + X_1(\vec{r}_\perp; t)) \\ F_2(x, \vec{r}_\perp; t) &= \tanh(x + d + X_2(\vec{r}_\perp; t)) \end{aligned} \quad (3.5)$$

The fields $X_{1,2}(\vec{r}_\perp; t)$ are determined by the boundary magnitudes of the field $\rho^{(cl)}(x = \pm d)$

$$\begin{aligned} \rho_{s1}(\vec{r}_\perp; t) &= \rho_1^{(cl)}(x = d, \vec{r}_\perp; t) = \sqrt{n_0} \tanh(X_1(\vec{r}_\perp; t)) \\ \rho_{s2}(\vec{r}_\perp; t) &= \rho_2^{(cl)}(x = -d, \vec{r}_\perp; t) = \sqrt{n_0} \tanh(X_2(\vec{r}_\perp; t)) \end{aligned} \quad (3.6)$$

Due to the large value of the external potential U_0 (or $\varkappa \gg 1$) the essential contribution into the integral for Z is given by the fields with the small values of the modulus at the boundary $|\psi_{s1,2}(\vec{r}_\perp; t)| \ll \sqrt{n_0}$. This means that the fields $X_{1,2}$ can be considered as the fields with the small values $|X_{1,2}| \ll 1$ and, thus, these fields can be expressed in the simple form in terms of the boundary values of the fields $\rho_{s1,2}(\vec{r}_\perp; t)$

$$X_{1,2}(\vec{r}_\perp; t) = \frac{1}{\sqrt{n_0}} \rho_{s1,2}(\vec{r}_\perp; t)$$

IV. DENSITY FLUCTUATIONS

For the consideration of the fluctuations of the modulus, the field ρ is represented as

$$\rho = \rho^{(cl)} + \sqrt{n_0} \delta\rho \quad (4.1)$$

The boundary conditions are taken into account in the definition of the field $\rho^{(cl)}$ and, therefore, the fluctuations of the modulus $\sqrt{n_0} \delta\rho$ should obey the zero boundary conditions

$$\delta\rho_1(x = d, \vec{r}_\perp; t) = \delta\rho_2(x = -d, \vec{r}_\perp; t) = 0 \quad (4.2)$$

Within the framework of the saddle-point approximation the inequality $|\delta\rho| \ll \rho^{(cl)}/\sqrt{n_0} = F$ should hold for. Substituting (4.1) into the action (2.6) and taking into account the inequality $|\delta\rho| \ll \rho^{(cl)}/\sqrt{n_0} = F$, we obtain

$$S_{vol} = S_{vol}^{(0)} + \delta S_{vol} \quad (4.3)$$

where

$$S_{vol}^{(0)} = n_0 \xi^3 \oint dt \int_{U(x)=0} dx \int d^2 r_\perp \left\{ -F^2 (\partial_t \varphi) + F^2 - \frac{1}{2} F^4 - \frac{1}{2} (\vec{\nabla} F)^2 \right\} \quad (4.4)$$

$$\delta S_{vol} = n_0 \xi^3 \oint dt \int_{U(x)=0} dx \int d^2 r_\perp \left\{ -2F \delta\rho (\partial_t \varphi) - \frac{1}{2} F^2 (\vec{\nabla} \varphi)^2 + \delta\rho \hat{H}_1 \delta\rho + \frac{1}{2} \delta\rho \vec{\nabla}_\perp^2 \delta\rho \right\} \quad (4.5)$$

where the function F is obtained by Eq. (3.5) and the operator $\hat{H}_1$ reads

$$\hat{H}_1 = -\frac{1}{2} \nabla_x^2 - 1 + 3F^2 \quad (4.6)$$

The spectrum of the operator $\hat{H}_1$ can be found [19] and has the form

$$\lambda_0 = 0, \quad \lambda_1 = \frac{3}{2}, \quad \lambda_k = 2 + \frac{1}{2} k^2 \quad (4.7)$$

Note that the zero mode self-energy λ_0 of operator $\hat{H}_1$ should be eliminated due to zero boundary conditions (4.2). At the same time the second discrete self-energy λ_1 and the self-energies of the continuous spectrum λ_k are not changed essentially by boundary conditions (4.2). In Eqs. (4.4), (4.5) the fluctuations of modulus $\delta\rho$ are taken into account within an accuracy to second order due to inequality $\delta\rho \ll F$ and the phase is taken into account exactly. The possibility of the consideration of the fluctuations $\delta\rho$ within an accuracy to second order results from the gap character of the spectrum of these fluctuations, in contrast to the spectrum of the fluctuations of the phase which is gapless.

The contribution of the density fluctuations into the generation functional Z is determined by action δS_{vol} and has the form of the Gaussian integral over $\delta\rho$. The integration over $\delta\rho$ in Z can be performed and we obtain

$$Z = \int DX D\varphi e^{i(S_{vol}^{(0)} + S^{(\varphi)} + S_{surf} + S_j)} \quad (4.8)$$

$$S^{(\varphi)} = n_0 \xi^3 \oint dt \int d^3 \vec{r} d^3 \vec{r}' \left\{ (F(\partial_t \varphi))_{\vec{r}} \left(\hat{H}_1 - \frac{1}{2} \vec{\nabla}_\perp^2 \right)_{\vec{r}, \vec{r}'}^{-1} (F(\partial_t \varphi))_{\vec{r}'} - \frac{1}{2} F^2 (\vec{\nabla} \varphi)^2_{\vec{r}} \delta[\vec{r} - \vec{r}'] \right\} \quad (4.9)$$

The evolution of the difference between the phases of the condensates in the right-hand and the left-hand sides of the bulk is a slow process. The inverse characteristic time of this process is proportional to the tunneling amplitude $\gamma = \frac{1}{2} n_0 (\kappa \sinh(2\kappa d))^{-1}$ and has the small value. Therefore, the momenta $\vec{k}_\perp$ of the fluctuations corresponding to these fluctuation frequencies should have small value $k_\perp \xi \ll 1$ (or in our system of units where $\xi = 1$, $k_\perp \ll 1$). As

a result, due to the gap character of the self-energies of operator $\hat{H}_1$ in the case of zero boundary conditions for $\delta\rho$ in the direction $\vec{r}_\perp$ we can neglect the term $\frac{1}{2}\vec{\nabla}_\perp^2$ compared with $\hat{H}_1$ in the first summand in the action (4.9).

The operator $\hat{H}_1$ can be represented in the factorized form

$$\hat{H}_1 = \frac{1}{2}\hat{L}_1^+\hat{L}_1 \quad (4.10)$$

where the operators $\hat{L}_1, \hat{L}_1^+$ have the form

$$\begin{aligned} \hat{L}_1 &= \nabla_x + 2F \\ \hat{L}_1^+ &= -\nabla_x + 2F \end{aligned} \quad (4.11)$$

Due to the possibility to represent $\hat{H}_1$ in the factorized form and to neglect $\vec{\nabla}_\perp^2$ in the first summand of Eq. (4.9), the action $S^{(\varphi)}$ can be written as

$$S^{(\varphi)} = n_0\xi^3 \oint dt \int_{U(x)=0} dx \int d^2\vec{r}_\perp \left\{ 2 \left[\left(\hat{L}_1^+ \right)^{-1} (F\partial_t\varphi) \right]^2 - \frac{1}{2}F^2 \left(\vec{\nabla}_\perp\varphi \right)^2 \right\} \quad (4.12)$$

After the integration by parts over x in the second term of the action $S^{(\varphi)}$ (4.12) this action can be transformed to the sum of two terms

$$S^{(\varphi)} = S_{vol}^{(\varphi)} + \tilde{S}_{surf}^{(\varphi)} \quad (4.13)$$

where

$$S_{vol}^{(\varphi)} = n_0\xi^3 \oint dt \int_{U(x)=0} dx \int d^2\vec{r}_\perp \left\{ 2 \left[\left(\hat{L}_1^+ \right)^{-1} (F\partial_t\varphi) \right]^2 - (F\varphi) \hat{H}_2 (F\varphi) - \frac{1}{2}F^2 \left(\vec{\nabla}_\perp\varphi \right)^2 \right\} \quad (4.14)$$

$$\tilde{S}_{surf}^{(\varphi)} = -\frac{1}{2}n_0\xi^3 \oint dt \int d^2\vec{r}_\perp \left\{ [F^2\varphi_1 (\nabla_x\varphi_1)] \Big|_{x=d}^{x=L} + [F^2\varphi_2 (\nabla_x\varphi_2)] \Big|_{x=-L}^{x=-d} \right\} \quad (4.15)$$

$$\hat{H}_2 = -\frac{1}{2}\nabla_x^2 - 1 + F^2 \quad (4.16)$$

The term $\tilde{S}_{surf}^{(\varphi)}$ of the action $S^{(\varphi)}$ is a surface one. The region of the integration in this surface term is the surface of the right-hand and the left-hand sides of the bulk.

Note that the operator $\hat{H}_2$ can be represented in the factorized form as well as operator $\hat{H}_1$

$$\hat{H}_2 = \frac{1}{2}\hat{L}_2^+\hat{L}_2 \quad (4.17)$$

where

$$\begin{aligned} \hat{L}_2 &= -\nabla_x + \frac{(\nabla_x F)}{F} \\ \hat{L}_2^+ &= \nabla_x + \frac{(\nabla_x F)}{F} \end{aligned} \quad (4.18)$$

It can easily be obtained that the operators $\hat{L}_1$ and $\hat{L}_2$ obey the following equations

$$\hat{L}_1\hat{H}_2\hat{L}_1^+ = \frac{1}{2}\hat{L}_1\hat{L}_2^+\hat{L}_2\hat{L}_1^+ = \frac{1}{2}\nabla_x (\nabla_x^2 - 4) \nabla_x \quad (4.19)$$

$$\hat{L}_1\hat{L}_1^+ = 2\hat{H}_2 + 4 \quad (4.20)$$

V. THE SURFACE TERMS OF THE ACTION

The part of the action $S_{vol}^{(0)}$ which consists of the modulus terms alone and does not have the phase fluctuations can be transformed to the surface form. This can easily be seen by the substitution of equation (3.5) for F into the action $S_{vol}^{(0)}$ (4.4). Taking into account the smallness of X ($X \ll 1$), we obtain

$$\begin{aligned} & n_0 \xi^3 \oint dt \int_{U(x)=0} dx \int d^2 r_{\perp} \left\{ F^2 - \frac{1}{2} F^4 - \frac{1}{2} (\nabla_x F)^2 - \frac{1}{2} (\vec{\nabla}_{\perp} F)^2 \right\} \\ &= n_0 \xi^3 \oint dt \int d^2 r_{\perp} \left\{ X - \frac{1}{3} (\vec{\nabla}_{\perp} X)^2 \right\} \end{aligned} \quad (5.1)$$

Note that taking into account the smallness of X and assuming the slowness of X compared with the scale ξ in the $\vec{r}_{\perp}$ space (the reason of this assumption has been discussed above), we can neglect the term $(\vec{\nabla}_{\perp} X)^2$.

The term of the action $S_{vol}^{(0)}$ depending on the fluctuation of the phase can be transformed to the surface form in the assumption of the slowness of the field φ compared with the scale ξ .

$$\begin{aligned} & n_0 \xi^3 \oint dt \int_{U(x)=0} dx \int d^2 r_{\perp} \{ F^2 (\partial_t \varphi) \} = \\ &= n_0 \xi^3 \oint dt \int_{U(x)=0} dx \int d^2 r_{\perp} \{ (F^2 - 1) (\partial_t \varphi) \} = \\ &= n_0 \xi^3 \oint dt \int d^2 r_{\perp} \{ X (\partial_t \varphi_s) \} \end{aligned} \quad (5.2)$$

As a result, the sum of $S_{vol}^{(0)}$ and S_{surf} is given by

$$S_{vol}^{(0)} + S_{surf} = n_0 \xi^3 \oint dt \int d^2 r_{\perp} \left\{ X - \frac{\varkappa}{2} X^2 - X (\partial_t \varphi_s) + \frac{\varkappa}{\sinh(2\varkappa d)} X^2 \cos(\Delta \varphi_s) \right\} \quad (5.3)$$

where $\Delta \varphi_s = \varphi_{s1} - \varphi_{s2}$ is the difference between the magnitudes of the phase fields at the boundary of the region of the nonzero external potential.

Due to the large value of $n_0 \xi^3 \gg 1$ the saddle-point approximation for the integral over X in the generation functional can be used. As can readily be seen, the saddle-point value of X is equal to X_0

$$X_0 = \frac{1}{\varkappa} \quad (5.4)$$

After integrating Z over the fluctuations of X near the saddle-point value X_0 and taking into account (4), we obtain the surface part of the action in the form

$$\tilde{S}_{surf}^{(\varphi)} = n_0 \xi^3 \oint dt \int d^2 r_{\perp} \left\{ -\frac{1}{2\varkappa} (\partial_t \varphi_s)^2 + \frac{1}{\varkappa \sinh(2\varkappa d)} \cos(\Delta \varphi_s) \right\} \quad (5.5)$$

Combining two surface terms $\tilde{S}_{surf}^{(\varphi)}$ (5.15) and $\tilde{S}_{surf}^{(\varphi)}$ (5), we obtain the surface term in the form

$$S_{surf}^{(\varphi)} = \tilde{S}_{surf}^{(\varphi)} + \tilde{S}_{surf}^{(\varphi)} \quad (5.6)$$

VI. THE EFFECTIVE ACTION FOR THE PHASE

Due to the smallness of the fluctuations of X we replace X with the saddle point value X_0 in the field F and denote this field as $F^{(0)}$ where

$$F^{(0)}(x) = \tanh\left(\frac{|x \mp d|}{\xi} + X_0\right) \quad (6.1)$$

The function $F^{(0)}$ is substituted in the action $S_{vol}^{(\varphi)}$ defined by Eq. (4.13) instead of F . The same procedure refers also to the operators $\hat{L}_1^+$, and $\hat{L}_1$. After this replacement the bulk part of the action $S_{vol}^{(\varphi)}$ can be written as

$$S_{vol}^{(\varphi)} = n_0 \xi^3 \oint dt \int_{U(x)=0} dx \int d^2 r_{\perp} \left\{ \begin{aligned} & 2 \left[\left(\hat{L}_1^{(0)+} \right)^{-1} (F^{(0)} \partial_t \varphi) \right]^2 - \\ & - (F^{(0)} \varphi) \hat{H}_2^{(0)} (F^{(0)} \varphi) - \frac{1}{2} (F^{(0)})^2 (\vec{\nabla}_{\perp} \varphi)^2 \end{aligned} \right\} \quad (6.2)$$

where the operators $\hat{L}_1^{(0)+}$ and $\hat{H}_2^{(0)}$ have the form of $\hat{L}_1^+$ and $\hat{H}_2$ with the substitution $F \rightarrow F^{(0)}$. Since the action $S_{vol}^{(\varphi)}$ has such form, that it is convenient to introduce the field η instead of the field φ

$$\eta = \left(\hat{L}_1^{(0)+} \right)^{-1} (F^{(0)} \varphi) \quad (6.3)$$

The transformation yields

$$S_{vol}^{(\varphi)} = n_0 \xi^3 \oint dt \int_{U(x)=0} dx \int d^2 r_{\perp} \left\{ \begin{aligned} & 2 (\partial_t \eta)^2 - \eta \left(\hat{L}_1^{(0)} \hat{H}_2^{(0)} \hat{L}_1^{(0)+} \right) \eta - \\ & - \frac{1}{2} (\vec{\nabla}_{\perp} \eta) \left(\hat{L}_1^{(0)} \hat{L}_1^{(0)+} \right) (\vec{\nabla}_{\perp} \eta) \end{aligned} \right\} \quad (6.4)$$

Using (4.19) and (4.20), the action $S_{vol}^{(\varphi)}$ can be written as

$$S_{vol}^{(\varphi)} = 2n_0 \xi^3 \oint dt \int_{U(x)=0} dx \int d^2 r_{\perp} \left\{ \begin{aligned} & (\partial_t \eta)^2 - \frac{1}{4} \eta \left[\nabla_x (\nabla_x^2 - 4) \nabla_x \right] \eta - \\ & - (\vec{\nabla}_{\perp} \eta) \left(\frac{1}{2} \hat{H}_2^{(0)} + 1 \right) (\vec{\nabla}_{\perp} \eta) \end{aligned} \right\} \quad (6.5)$$

In the case of the slow spatial variation of the field η along the x axis the action (6.5) can be transformed to

$$S_{vol}^{(\varphi)} = 2n_0 \xi^3 \oint dt \int_{U(x)=0} dx \int d^2 r_{\perp} \left\{ (\partial_t \eta)^2 + \eta (\nabla_x^2) \eta - \frac{1}{2} (\vec{\nabla}_{\perp} \eta) \left(1 + (F^{(0)})^2 \right) (\vec{\nabla}_{\perp} \eta) \right\} \quad (6.6)$$

The transition to the Fourier transformation in the $\vec{r}_{\perp}$ plane gives

$$S_{vol}^{(\varphi)} = 2n_0 \xi^3 \oint dt \int_{U(x)=0} dx \int \frac{d^2 k_{\perp}}{(2\pi)^2} \left\{ \eta_{-\vec{k}_{\perp}} \left[-\partial_t^2 + \nabla_x^2 - \vec{k}_{\perp}^2 - \frac{1}{2} \vec{k}_{\perp}^2 (\nabla_x F^{(0)}) \right] \eta_{\vec{k}_{\perp}} \right\} \quad (6.7)$$

The surface terms in the action $\tilde{S}_{surf}^{(\varphi)}$ (4.15) can easily be expressed in terms of the field η . Using the connection between the fields φ and η (6.3) having the form

$$\varphi = \frac{1}{F^{(0)}} \left(-\nabla_x + 2F^{(0)} \right) \eta \quad (6.8)$$

we express $\nabla_x \varphi$ in terms of η

$$\nabla_x \varphi = \frac{1}{F^{(0)}} \left(-\nabla_x + \frac{1 + (F^{(0)})^2}{F^{(0)}} \right) \nabla_x \eta \quad (6.9)$$

The characteristic frequency ω_0 of the phase time evolutions is proportional to the tunneling amplitude $\omega_0 \sim \gamma$, where the tunneling amplitude $\gamma = (\varkappa \sinh(2\varkappa d))^{-1}$. The tunneling amplitude γ is the small value being much smaller than $\frac{1}{\varkappa} = F^{(0)}(\pm d) \ll 1$, i.e., μ in the usual system of units. Thus, the characteristic frequency of the field η is small and, therefore, $\nabla_x \eta$ is small compared with η due to the connection $\omega \sim k_x$ on the mass surface for the small ω (Eq. 6.7). In this case Eq. (6.9) goes over into

$$\nabla_x \varphi = \frac{1 + (F^{(0)})^2}{(F^{(0)})^2} \nabla_x \eta \quad (6.10)$$

From Eqs. (6.8) and (6.10) due to the inequality $\omega_0 \sim k_x^{(0)} \sim \gamma \ll F^{(0)} (\pm d)$ ($k_x^{(0)}$ is the characteristic momentum corresponding to ω_0) we obtain

$$\varphi(d) = 2\eta(d) \quad (6.11)$$

$$\nabla_x \varphi(d) = \frac{1}{(F^{(0)}(d))^2} \nabla_x \eta(d)$$

$$\varphi(L) = 2\eta(L)$$

$$\nabla_x \varphi(L) = 2\nabla_x \eta(L)$$

From these equalities the surface terms in the Eq. (4.15) for $\tilde{S}_{surf}^{(\varphi)}$ can be written in terms of the η field in the form

$$\frac{1}{2} \left[\left(F^{(0)} \right)^2 \varphi \nabla_x \varphi \right] (d) = [\eta \nabla_x \eta] (d) \quad (6.12)$$

VII. THE INTEGRAL OVER THE PHASE

Using expressions (6.5-6.7) for action $S_{vol}^{(\varphi)}$, the integral over the field $\varphi(x, \vec{r}_\perp; t)$ with the fixed values $\varphi_s(\vec{r}_\perp; t)$ on the surface of the region of the nonzero external potential can be calculated. Note that the phases of the left-hand and the right-hand condensates are homogeneous in the $\vec{r}_\perp$ plane and do not depend on $\vec{r}_\perp$ due to the translational invariance of the system in these directions. Therefore, the surface phase fields $\varphi_s(\vec{r}_\perp; t)$ depending on $\vec{r}_\perp$ correspond to non-condensate particles and can be considered as perturbations. We analyse the relaxation of a difference in phases in zero order of these perturbations and do not consider them in this work, i.e., we put $\varphi_{s1}(t)$ and $\varphi_{s2}(t)$ depending on the time variable alone.

The action $S_{vol}^{(\varphi)}$ (6.5) has the Gaussian form. In this connection, in order to calculate the integral over the bulk components of the field η for the small ω and $\vec{k}$, we should find the solution of the equation

$$\left[\omega^2 + \nabla_x^2 - \vec{k}_\perp^2 + \frac{1}{2} \vec{k}_\perp^2 \left(\nabla_x F^{(0)} \right) \right] \eta = 0 \quad (7.1)$$

as a function of the boundary values of the field η and put this solution into the action $S_{vol}^{(\varphi)}$. We supposed that the boundary values of the phase field φ , i.e., φ_s and φ_L , are known. In the case of the boundary magnitudes of the field φ and, hence, of the field η independent on $\vec{r}_\perp$ we look for the solution of this equation for $\vec{k}_\perp = 0$, i.e.,

$$[\omega^2 + \nabla_x^2] \eta = 0 \quad (7.2)$$

As was mentioned above, to describe the relaxation, we use the Keldysh-Schwinger technique with the two-time contour [14], [15]. In this technique the field η has two components corresponding to two branches of the time contour (quadrangular representation). It is convenient to introduce the components of the field η as a sum and a difference of the two components corresponding to the upper and to the lower branches of the time contour (triangular representation) [14], [15]. In Eqs. (7.1), (7.2) the frequency ω is a complex quantity $\omega \rightarrow \omega + i\Gamma$. The imaginary part Γ is the relaxation frequency of the excitations in the homogeneous Bose gas. The value Γ depends on ω and for the small ω this value is much smaller than ω , i.e., $\omega \gg \Gamma$ [20]. Thus, the equation for the first component of the field η has the frequency ω in the form of $\omega \rightarrow \omega^* = \omega - i\Gamma$ and the equation for the second one has the frequency in the form $\omega \rightarrow \omega = \omega + i\Gamma$.

The Keldysh Green function of this equation is denoted as $\hat{D}(x, 0)$. We use the "triangular" representation. The solution of Eq. (7.2) with the fixed boundary conditions is

$$\eta(x) = \hat{D}(x, 0) \left(\hat{D}(0, 0) \right)^{-1} \eta(0) \quad (7.3)$$

the coordinates $x=\pm d$ of the boundary we denote as 0. The Green function $\hat{D}(x, 0)$ in the "triangular" representation reads

$$\hat{D} = \begin{pmatrix} 0 & D_A \\ D_R & D_K \end{pmatrix} \quad (7.4)$$

where D_R and D_A are the retarded and advanced Green functions respectively, and D_K is the kinetic Green function. For the equilibrium state, the Green function D_K is connected with the functions D_R and D_A by the equation

$$D_K = \coth\left(\frac{\omega}{2T}\right) (D_R - D_A) \quad (7.5)$$

The Green functions D_R and D_A have the form

$$\begin{aligned} D_R(x, 0) &= \int \frac{dp_x}{2\pi} \frac{e^{ixp_x}}{p_x^2 - (\omega + i\Gamma)^2} = \\ &= \theta(x) \frac{i}{2(\omega + i\Gamma)} e^{ix(\omega + i\Gamma)} - \theta(-x) \frac{i}{2(\omega + i\Gamma)} e^{-ix(\omega + i\Gamma)} \\ D_A(x, 0) &= \int \frac{dp_x}{2\pi} \frac{e^{ixp_x}}{p_x^2 - (\omega - i\Gamma)^2} = \\ &= \theta(-x) \frac{i}{2(\omega + i\Gamma)} e^{ix(\omega - i\Gamma)} - \theta(x) \frac{i}{2(\omega + i\Gamma)} e^{-ix(\omega + i\Gamma)} \end{aligned} \quad (7.6)$$

The inverse Green function $\hat{D}^{-1}$ in the "triangular" form is

$$\hat{D}^{-1} = \begin{pmatrix} -D_K D_R^{-1} D_A^{-1} & D_R^{-1} \\ D_A^{-1} & 0 \end{pmatrix} = \begin{pmatrix} \coth\left(\frac{\omega}{2T}\right) (D_R^{-1} - D_A^{-1}) & D_R^{-1} \\ D_A^{-1} & 0 \end{pmatrix} \quad (7.7)$$

The space derivatives of these Green functions give ($x>0$)

$$\begin{aligned} \nabla_x D_R(x, 0) &= -\frac{1}{2} e^{ix(\omega + i\Gamma)} \\ \nabla_x D_A(x, 0) &= -\frac{1}{2} e^{-ix(\omega - i\Gamma)} \end{aligned} \quad (7.8)$$

Using Eq. (7.3), we can represent the surface terms (5.12) and (4.15) of the action $\tilde{S}_{surf}^{(\varphi)}$ as

$$\begin{aligned} [\eta \nabla_x \eta](0) &= \\ &= \begin{pmatrix} \eta_1(x) & \eta_2(x) \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & \nabla_x D_A(x, 0) \\ \nabla_x D_R(x, 0) & \nabla_x D_K(x, 0) \end{pmatrix} \left(\hat{D}(0, 0) \right)^{-1} \begin{pmatrix} \eta_1(0) \\ \eta_2(0) \end{pmatrix} \end{aligned} \quad (7.9)$$

$$\begin{pmatrix} 0 & \nabla_x D_A(0) \\ \nabla_x D_R(0) & \nabla_x D_K(0) \end{pmatrix} = -\frac{1}{2} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad (7.10)$$

$$\begin{aligned} \left(\hat{D}(0, 0) \right)^{-1} &= \begin{pmatrix} \coth\left(\frac{\omega}{2T}\right) (D_R^{-1}(0, 0) - D_A^{-1}(0, 0)) & D_R^{-1}(0, 0) \\ D_A^{-1}(0, 0) & 0 \end{pmatrix} = \\ &= 2 \begin{pmatrix} -2i\omega \coth\left(\frac{\omega}{2T}\right) & -i\omega \\ i\omega & 0 \end{pmatrix} \end{aligned} \quad (7.11)$$

Thus, we have

$$[\eta \nabla_x \eta](0) = \begin{pmatrix} \eta_1(0) & \eta_2(0) \end{pmatrix} \begin{pmatrix} -2i\omega \coth\left(\frac{\omega}{2T}\right) & -i\omega \\ i\omega & 0 \end{pmatrix} \begin{pmatrix} \eta_1(0) \\ \eta_2(0) \end{pmatrix} \quad (7.12)$$

Substituting (7.12) into (4.15), we obtain the following expression for the action $\tilde{S}_{surf}^{(\varphi)}$ in the "triangle" representation

$$\tilde{S}_{surf}^{(\varphi)} = -\frac{1}{8}n_0\xi^3 S_{\perp} \int dt \left\{ \left(\begin{smallmatrix} \phi_s \\ \Phi_s \end{smallmatrix} \right) \left(\hat{D}(0,0) \right)^{-1} \left(\begin{smallmatrix} \phi_s \\ \Phi_s \end{smallmatrix} \right) \right\} \quad (7.13)$$

In the time representation the frequency ω is replaced by the time derivative $i\partial_t$. The fields Φ and ϕ are defined by the equalities

$$\begin{aligned} \Phi &= \frac{1}{2}(\varphi_1 + \varphi_2) \\ \phi &= \frac{1}{2}(\varphi_1 - \varphi_2) \end{aligned} \quad (7.14)$$

where φ_1 and φ_2 are the phase fields determined on the upper and the lower brunches of the time contour, respectively.

As a result, the action $S_{surf}^{(\varphi)}$ as an effective action S_{eff} which describes the relaxation of the difference between the phases at the boundary of the region of the nonzero external potential can be written in the "triangular" form as

$$\begin{aligned} S_{eff} &= S_{surf}^{(\varphi)} = \\ &= n_0\xi^3 S_{\perp} \int dt \left\{ \sum_{\alpha=1,2} \left(\begin{smallmatrix} \phi_{s\alpha} \\ \Phi_{s\alpha} \end{smallmatrix} \right) \left[\left(\hat{G}_s^{(0)} \right)^{-1} - \frac{1}{8} \left(\hat{D}(0,0) \right)^{-1} \right] \left(\begin{smallmatrix} \phi_{s\alpha} \\ \Phi_{s\alpha} \end{smallmatrix} \right) - \right. \\ &\quad \left. - (V[\Delta\Phi_s + \Delta\phi_s] - V[\Delta\Phi_s - \Delta\phi_s]) \right\} \end{aligned} \quad (7.15)$$

where

$$\left(\hat{G}_s^{(0)} \right)^{-1} = \frac{1}{2\kappa} \begin{pmatrix} 0 & \omega^2 \\ \omega^2 & 0 \end{pmatrix} \quad (7.16)$$

$$V(\varphi_s) = -\frac{1}{\kappa \sinh(2\kappa d)} \cos(\varphi_s) \quad (7.17)$$

The index $\alpha = 1, 2$ in Eq. (7.15) corresponds to the right-hand or the left-hand side of the bulk and $\Delta\Phi_s = \Phi_{s1} - \Phi_{s2}$, $\Delta\phi_s = \phi_{s1} - \phi_{s2}$. The value $S_{\perp}$ is the area of the section of the bulk along the y-z plane.

VIII. THE DESCRIPTION OF THE RELAXATION

The effective action (7.15) describes the relaxation of the difference between the phases at the right and the left boundaries of the region of the nonzero external potential. The generation functional Z can be written in the form

$$Z = \int D\phi D\Phi e^{i(S_{eff} + S_j)} \quad (8.1)$$

To simplify the consideration, we suppose that the temperature T is much larger than the characteristic frequency $\omega_0 \sim \gamma = (\kappa \sinh(2\kappa d))^{-1}$. In this assumption, firstly, the term $\coth(\frac{\omega}{2T})$ in $\left(\hat{D}(0,0) \right)^{-1}$ (7.11) can be written as $\frac{2T}{\omega}$. Secondly, the component of the phase field ϕ describing the quantum fluctuations is much smaller than the component Φ ($\phi \gg \Phi$). This approximation is conditioned by the large value of the parameter $n_0\xi^3 S_{\perp} \gg 1$ (the system of units is such that the scale of the length is defined by the value ξ). Due to the smallness of the field ϕ compared with the field Φ it is possible to expand the nonlinear terms $V(\Delta\Phi_s \pm \Delta\phi_s)$ in a series in $\Delta\phi_s$ to first order. Thus, we obtain

$$S_{eff} = n_0\xi^3 S_{\perp} \int dt \left\{ \sum_{\alpha=1,2} \phi_{s\alpha} \left[\frac{1}{2\kappa} \omega^2 + \frac{i}{4} \omega \right] \Phi_{s\alpha} + \Phi_{s\alpha} \left[\frac{1}{2\kappa} \omega^2 - \frac{i}{4} \omega \right] \phi_{s\alpha} + i\phi_{s\alpha} T \phi_{s\alpha} - \right. \\ \left. - 2(\Delta\phi_s) V'[\Delta\Phi_s] \right\} \quad (8.2)$$

The term $V'(\Delta\Phi_s)$ is the functional derivative of the functional $V[\Delta\Phi_s]$, i.e., $V'[\Delta\Phi_s] = \delta V[\Delta\Phi_s] / \delta(\Delta\Phi_s)$.

We introduce the half-sum and the half-difference of the fields $\phi_{s\alpha}$ and $\Phi_{s\alpha}$

$$\begin{aligned}\tilde{\Phi} &= \frac{1}{2} (\Phi_{s1} + \Phi_{s2}) ; & \tilde{\phi} &= \frac{1}{2} (\phi_{s1} + \phi_{s2}) \\ (\Delta\Phi) &= \frac{1}{2} (\Phi_{s1} - \Phi_{s2}) ; & (\Delta\phi) &= \frac{1}{2} (\phi_{s1} - \phi_{s2})\end{aligned}\tag{8.3}$$

After the integration of the generation potential Z over the fields $\tilde{\Phi}$ and $\tilde{\phi}$ we obtain

$$Z = \int D(\Delta\phi) D(\Delta\Phi) e^{i(\Delta S_{eff} + S_j)}\tag{8.4}$$

where

$$\Delta S_{eff} = n_0 \xi^3 S_{\perp} \int dt \left\{ (\Delta\phi) \left[\frac{1}{2\kappa} \omega^2 + \frac{i}{4} \omega \right] (\Delta\Phi) + (\Delta\Phi) \left[\frac{1}{2\kappa} \omega^2 - \frac{i}{4} \omega \right] (\Delta\phi) + \right. \\ \left. + i (\Delta\phi) T(\Delta\Phi) - 2 (\Delta\phi) V'[\Delta\Phi] \right\}\tag{8.5}$$

The fields $\Delta\Phi$ and $\Delta\phi$ are represented as a sum of the parts changing slowly and fast in time. We denote the fast parts of $\Delta\Phi$ and $\Delta\phi$ as $\delta\Phi$ and $\delta\phi$ and the slow parts as Φ and ϕ . Thus

$$\begin{aligned}\Delta\Phi &\rightarrow \Phi + \delta\Phi \\ \Delta\phi &\rightarrow \phi + \delta\phi\end{aligned}\tag{8.6}$$

The criterion of separating the fields into the fast and slow ones is the magnitude of the frequency. If the frequency of the field $\omega \gtrsim \omega_c$ where $\omega_c = 1/\kappa$, the field is fast, correspondingly, if the frequency $\omega \ll \omega_c$, the field is slow. This gives

$$S_{eff} = S_{eff}^{(slow)} + S_{eff}^{(fast)}\tag{8.7}$$

$$S_{eff}^{(slow)} = n_0 \xi^3 S_{\perp} \int dt \left\{ \phi \left[\frac{1}{2\kappa} \omega^2 + \frac{i}{4} \omega \right] \Phi + \Phi \left[\frac{1}{2\kappa} \omega^2 - \frac{i}{4} \omega \right] \phi + \right. \\ \left. + i \phi T \Phi - 2 \phi V'[\Phi] \right\}\tag{8.8}$$

$$S_{eff}^{(fast)} = n_0 \xi^3 S_{\perp} \int dt \left\{ (\delta\phi) \left[\frac{1}{2\kappa} \omega^2 + \frac{i}{4} \omega - V''[\Phi] \right] (\delta\Phi) + \right. \\ \left. (\delta\Phi) \left[\frac{1}{2\kappa} \omega^2 - \frac{i}{4} \omega - V''[\Phi] \right] (\delta\phi) + i (\delta\phi) T(\delta\phi) \right\}\tag{8.9}$$

The integration over $\delta\phi$ yields

$$S_{\Phi}^{(fast)} = n_0 \xi^3 S_{\perp} \int dt \left\{ i \frac{1}{16T} [\hat{L}(\delta\Phi)]^2 \right\}\tag{8.10}$$

where the operator $\hat{L}$ is

$$\hat{L} = \frac{1}{2\kappa} \partial_t^2 + \frac{1}{4} \partial_t + V''(\Phi)\tag{8.11}$$

Note that for the slow frequencies $\omega \ll 1$, the term $\frac{1}{2\kappa} \partial_t^2$ can be neglected compared with the term $\frac{1}{4} \partial_t$.

For the generation potential, we obtain

$$Z = Z_{slow} Z_{fast}\tag{8.12}$$

where

$$Z_{slow} = \int D\phi D\Phi e^{i(S_{eff}^{(slow)} + S_j^{(slow)})}\tag{8.13}$$

$$Z_{fast} = \int D(\delta\Phi) e^{i(S_{\Phi}^{(fast)} + S_j^{(fast)})}\tag{8.14}$$

Due to the form of the action $S_{\Phi}^{(fast)}$ (8.10) the generation potential Z_{fast} has the Gaussian form and can be calculated

$$Z_{fast} = C_1 \det_1 [\hat{L}^{-1}] \quad (8.15)$$

where the constant C has the form

$$C_1 = \int D(\delta\Phi) \exp \left\{ - \int dt \left\{ \frac{1}{16T} (\delta\Phi)^2 \right\} \right\}$$

and can be included into a measure of the functional integral Z_{slow} . The index 1 in $\det_1$ denotes the product of the Fourier components corresponding to the fast frequencies. The value $\det_1 [\hat{L}^{-1}]$ can be calculated by the following way

$$\begin{aligned} \det_1 [\hat{L}^{-1}] &= e^{Sp'' Ln[\hat{L}^{-1}]} = C_2 \exp \left\{ Sp'' Ln \left[\left(G_R^{(0)} \right) V''(\Phi) \right] \right\} = \\ &= C_2 \exp \left\{ -Sp' Ln \left[\left(G_R^{(0)} \right) V''(\Phi) \right] \right\} \end{aligned} \quad (8.16)$$

The symbol Sp'' indicates the sum over the fast frequencies and the symbol Sp' indicates the sum over the slow frequencies. The inessential c-number constant C_2 is equal to

$$C_2 = \exp \left\{ -Sp'' Ln \left[\left(G_R^{(0)} \right)^{-1} \right] \right\}$$

and can be included into the measure of the Z_{slow} . The value $Sp' Ln \left[\left(G_R^{(0)} \right) V''(\Phi) \right]$ is calculated as

$$Sp' Ln \left[\left(G_R^{(0)} \right) V''(\Phi) \right] = V''(\Phi(t)) \int_{-\frac{1}{4}}^{\frac{1}{4}} \frac{d\omega}{2\pi} \frac{1}{\frac{i}{4}(\omega + i\Gamma)} = -4\theta(0) V''(\Phi) = -2V''(\Phi)$$

Hence, the expression for the generation potential has the form

$$Z_{fast} = C_1 \det_1 [\hat{L}^{-1}] = \exp \left\{ 2 \int dt V''(\Phi) \right\} \quad (8.17)$$

This equation gives

$$Z = Z_{slow} Z_{fast} = \int D\phi D\Phi e^{i(S_{eff}^{(slow)} + S_j^{(slow)} - 2i \int dt V''(\Phi))} \quad (8.18)$$

The action $S_{eff}^{(fast)}$ has form (8.9). Integrating Z (8.18) over the field ϕ , we obtain

$$Z = \int D\Phi e^{-\left(S_{slow}^{(\Phi)} - 2 \int dt V''(\Phi) - i S_j^{(slow)} \right)} \quad (8.19)$$

$$S_{slow}^{(\Phi)} = \int dt \left\{ \frac{1}{T} \left[\frac{1}{4} (\partial_t \Phi)^2 - V'(\Phi) \right]^2 - 2V''(\Phi) \right\} \quad (8.20)$$

Omitting the term $(\partial_t \Phi) V'(\Phi)$ which can be represented as a total derivative, we obtain

$$S_{slow}^{(\Phi)} = \frac{1}{T} \int dt \left\{ \frac{1}{16} (\partial_t \Phi)^2 + (V'(\Phi))^2 - 2TV''(\Phi) \right\} \quad (8.21)$$

Note that the generation potential Z with action $S_{slow}^{(\Phi)}$ (8.21) has the form corresponding to the quantum mechanics with the "imaginary time" in which the role of the "imaginary time" plays the usual real time. This action corresponds to the action which is obtained for the classical random process describing by the Langeven equation with the white noise [21] and having the form of the supersymmetric quantum mechanics.

The Hamiltonian corresponding to the action $S_{slow}^{(\Phi)}$ (8.21) is

$$\hat{H} = -4T^2 \frac{d^2}{d\Phi^2} + (V'(\Phi))^2 - 2TV''(\Phi) \quad (8.22)$$

and the Schredinger equation corresponding to this Hamiltonian can be written as

$$-\partial_t \Psi = \hat{H} \Psi \quad (8.23)$$

The Hamiltonian $\hat{H}$ can be represented in the factorized form

$$\hat{H} = -(2T)^2 \left(\frac{d}{d\Phi} - A \right) \left(\frac{d}{d\Phi} + A \right) \quad (8.24)$$

where

$$A = \frac{V'(\Phi)}{2T} \quad (8.25)$$

The Hamiltonian $\hat{H}$ has the minimum eigenvalue equal to zero. The eigenwave function corresponding to the zero eigenvalue is

$$\Psi_0 = \exp \left\{ \int d\Phi A(\Phi) \right\} = \exp \left\{ -\frac{V(\Phi)}{2T} \right\} \quad (8.26)$$

This can easily be seen from the form of the Hamiltonian (8.24). Due to the form of the Hamiltonian (8.24) the Schredinger Eq. (8.23) can be modified into the form of the Fokker-Planck equation [18] by the transformation

$$\Psi \rightarrow \exp \left\{ -\int_0^\Phi d\Phi' A(\Phi') \right\} \Psi \quad (8.27)$$

In this case the eigenwave function (8.26) corresponds to the equilibrium distribution function for the field Φ which is formed as a result of the relaxation process.

This research was supported by the Russian Foundation for Basic Research and by the Netherlands Organization for Scientific Research (NWO).

REFERENCES

- [1] M. H. Anderson, J. H. Ensher, M. B. Matthews, C. E. Weiman, and E. A. Cornell, Science 269, 198, (1995); K. B. Davis et al., Phys. Rev. Lett. 75, 3969, (1995); C. C. Bradley, C. A. Sackett, J. T. Tolleth, and R. G. Hulet, Phys. Rev. Lett. 75, 1687, (1995).
- [2] M. R. Andrews, C. G. Townsend, H. J. Meisner, Science 275, 637, (1997).
- [3] A. Rohrl, Naraschewski, A. Schenzle, and H. Wallis, Phys. Rev. Lett. 78, 4143, (1997)
- [4] D. S. Hall, M. R. Matthews, C. E. Weiman, and E. A. Cornell, Phys. Rev. Lett. 81, 1543, (1998).
- [5] J. Javanainen, Phys. Rev. Lett. 57, 3164, (1986).
- [6] F. Dalfovo, L. P. Pitaevskii, and S. Stringari, Phys. Rev. A 54, 4213, (1996).
- [7] A. Smerzi, S. Fantoni, S. Giovanazzi, and S. R. Shenoi, Phys. Rev. Lett. 79, 4950, (1997).
- [8] I. Zapata, F. Sols, and Leggett, Phys. Rev. A 57, R28, (1998).
- [9] B. D. Josephson, Phys. Lett. 1, 251, (1962).
- [10] V. Ambegaokar, U. Eckern, and G. Schen, Phys. Rev. Lett. 48, 1745, (1982), U. Eckern, and G. Schen, V. Ambegaokar, Phys. Rev. B 30, 6419, (1984).
- [11] F. Meier, and W. Zwerger, cond-mat/ 9904147.
- [12] S. T. Belyaev, JETP, 34, 417, (1958).
- [13] A. A. Abrikosov, L. P. Gor'kov, I. E. Dzyaloshinskii, "Methods of Quantum Field Theory in Statistical Physics", Moscow, 1998.
- [14] L. V. Keldysh, JETP, 47, 1515, (1964).
- [15] J. Schwinger, J. Math. Phys. 2, 407, (1961).
- [16] D. L. Kovrizhin, Phys. Lett. A 287, 392 (2001).
- [17] R. P. Feynman, A. R. Hibbs, "Quantum Mechanics and Path Integrals", N.Y., 1965.

- [18] H. Risken, Z. Phys., 251,231, (1972); K. Kawasaki, Progr. Theor. Phys. 51, 1064, (1974); R. Graham Z. Phys. B40, 149, (1980).
- [19] L. D. Landau, E. M. Lifshits, "Quantum Mechanics", M., 1974.
- [20] I. M. Khalatnikov, JETP 23, 21 (1952).
- [21] M. V. Feigelman, A. M. Tselik, JETP, 83, (1982), 1430.